

LETTER TO THE EDITOR

Genital necrosis with cutaneous thrombosis after COVID-19 mRNA vaccination

Editor

Thrombosis is a rare complication of COVID-19 vaccines that typically affects cerebral and visceral vessels.^{1–4} However, skin involvement is largely unknown.^{5,6} Here, we describe a case of genital necrosis associated with cutaneous thrombosis following COVID-19 vaccination.

An 84-year-old Japanese woman presented to our department with a three-day history of genital necrosis. She had received her first dose of Pfizer–BioNTech (New York, NY, USA; Mainz, Germany) BNT162b2 mRNA COVID-19 vaccine 26 days before admission. Nine days after the vaccination, she developed increasing pain in her genital region. She denied any trauma or precipitating event. Her medical history was significant for deep vein thrombosis after orthopaedic surgery, for which she had been receiving edoxaban over the past three years. She had no other risk factors for thrombosis.

On admission, she was well but febrile to 37.5°C. Dermatological examination revealed extensive necrosis with surrounding purpura that involved the mons pubis, labia majora and perineum (Fig. 1a). Laboratory investigations showed a leukocytosis ($15.9 \times 10^9/L$) with a left shift. The platelet count was slightly elevated ($359 \times 10^9/L$). The coagulation profile was unremarkable. Biochemical parameters were within the normal range except for an elevated C-reactive protein (11.6 mg/dL, normal <0.3 mg/dL). A thrombophilia screen—including anti-thrombin, protein C, protein S, lupus anticoagulant, anti-cardiolipin antibodies and anti- β -2-glycoprotein-1 antibodies—was unremarkable. Serological tests for rheumatoid factor, antinuclear antibody and anti-neutrophil cytoplasmic antibodies were all negative. Pelvic CT was performed to show subcutaneous fat stranding without fascial thickening. No haemorrhage or hematoma was noted. CT angiography detected no evidence of thrombosis. Skin biopsy showed epidermal necrosis, scattered neutrophils and lymphocytes in the dermis, and thrombotic occlusion of dermal vessels with mild perivascular infiltration (Fig. 1b,c). Immunohistochemistry revealed that the thrombi were positive for CD61, a platelet-specific marker (Fig. 1d,e). Based on the clinical and histopathological findings, a diagnosis of cutaneous necrosis with platelet thrombi formation and secondary infection was made. Treatment was started with




Figure 1 Skin lesions and histopathological findings at the time of admission. (a) Extensive necrosis with purpura in the genital region. (b,c) Histopathology showing epidermal necrosis and thrombotic occlusion of dermal vessels (haematoxylin-eosin stain, original magnification $\times 100$ [b] and $\times 200$ [c]). (d,e) Immunohistochemistry showing that the thrombi were positive for CD61 (original magnification $\times 100$ [d] and $\times 200$ [e]). Scale bar = 50 μ m (b,c).

ampicillin/sulbactam along with local wound care. Her fever, leukocytosis and genital pain resolved within the first week. The skin lesions also improved: more than 80% of the eschar had fallen off when she was discharged after one month of admission (Fig. 2a), and epithelization was almost completed another month later (Fig. 2b).

A small but increasing number of thrombotic events have been reported since the launch of mass vaccination campaigns against COVID-19. Adenovirus vector-based vaccines from AstraZeneca (Cambridge, UK) and Johnson & Johnson (Titusville, NJ, USA) are associated with severe thrombosis with thrombocytopenia, while mRNA-based vaccines from Pfizer–BioNTech and Moderna (Cambridge, MA, USA) are also associated with some thrombotic events, which do not always accompany thrombocytopenia.^{1–4} The exact pathogenesis remains unknown, but platelet activation is thought to be a key feature underlying these events.⁷ For both types of vaccines, thrombosis typically occurs in unusual locations such as cerebral and portal veins. However, only two cases of skin involvement have been reported, both of which manifested as local skin necrosis at injection sites.^{5,6} To the best of our knowledge, this is the first case of extensive skin necrosis after COVID-19 vaccination that




Figure 2 Improvement of the skin lesions. (a) Most of the eschar had fallen off one month after admission. (b) Epithelization was almost completed one month after discharge.

developed outside the injection site. Although the mechanism of platelet thrombi formation in genital skin is unclear, the short time interval between the vaccination and the onset of symptoms may indicate a causal relationship. Therefore, our case extends the range of cutaneous manifestations associated with thrombosis after COVID-19 vaccination. In addition, it should be noted that she developed thrombosis despite receiving edoxaban, which highlights the need to consider the possibility of thrombosis even in patients under anticoagulant therapy. Although rare, skin necrosis should be recognized as a possible manifestation of thrombosis associated with COVID-19 vaccination.

Acknowledgement

The patient in this manuscript has given written informed consent to publication of her case details.

Conflicts of interest

The authors have no conflict of interest to declare.

Funding source

None.

Data availability statement

The data presented in this manuscript are available from the corresponding author upon reasonable request.

A. Kuzumi, A. Yoshizaki,* K. Chiba, S. Mitsuo, K.M. Matsuda, Y. Norimatsu, K. Nagai, J. Omatsu, T. Miyake, S. Sato

Department of Dermatology, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

*Correspondence: A. Yoshizaki. E-mail: ayuyoshi@me.com

References

- Greinacher A, Thiele T, Warkentin TE, Weisser K, Kyrle PA, Eichinger S. Thrombotic thrombocytopenia after ChAdOx1 nCov-19 vaccination. *N Engl J Med* 2021; **384**: 2092–2101.
- Muir KL, Kallam A, Koepsell SA, Gundabolu K. Thrombotic Thrombocytopenia after Ad26.COV2.S Vaccination. *N Engl J Med* 2021; **384**: 1964–1965.
- Fan BE, Shen JY, Lim XR *et al*. Cerebral venous thrombosis post BNT162b2 mRNA SARS-CoV-2 vaccination: a black swan event. *Am J Hematol* 2021; **96**: E357–E361.
- Syed K, Chaudhary H, Donato A. Central venous sinus thrombosis with subarachnoid hemorrhage following an mRNA COVID-19 vaccination: are these reports merely co-incidental? *Am J Case Rep* 2021; **22**: e933397.
- Ramessur R, Saffar N, Czako B, Agarwal A, Batta K. Cutaneous thrombosis associated with skin necrosis following Oxford-AstraZeneca COVID-19 vaccination. *Clin Exp Dermatol* 2021; **46**: 1610–1612.
- Gruenstein D, Levitt J. Skin necrosis at both COVID-19 vaccine injection sites. *JAAD Case Rep* 2021; **15**: 67–68.
- Elrashdy F, Tambuwala MM, Hassan SS *et al*. Autoimmunity roots of the thrombotic events after COVID-19 vaccination. *Autoimmun Rev* 2021; **20**: 102941.

DOI: 10.1111/jdv.17837